

Search for Heavy Top-like Quark Pair Production in the Dilepton Final State in pp Collisions at $\sqrt{s} = 7 \text{ TeV}$

D. Barge, C. Campagnari, D. Kovalskyi, V. Krutelyov

University of California, Santa Barbara

W. Andrews, G. Cerati, D. Evans, F. Golf, I. Macneill, S. Padhi, Y. Tu, F. Würthwein, A. Yagil, J. Yoo

University of California, San Diego

L. Bauerdick, K. Burkett, I. Fisk, Y. Gao, O. Gutsche, B. Hooberman, S. Jindariani, J. Linacre, S. Tkaczyk

Fermi National Accelerator Laboratory, Batavia, Illinois

Approval

Exotica Meeting
24th January 2012



CMS PAPER EXO-11-050

Available on CMS information server

CMS AN -2011/509

DRAFT CMS Paper

The content of this note is intended for CMS internal use and distribution only


2012/01/17
Head Id: 95561
Archive Id: 95763P
Archive Date: 2012/01/17
Archive Tag: trunk

Search for Heavy Top-like Quark Pair Production in the
Dilepton Final State in pp Collisions at $\sqrt{s} = 7$ TeV

The CMS Collaboration


Abstract

The results of a search for pair production of a heavy top-like quark, t' , in the decay mode $t'\bar{t}' \rightarrow bW^+\bar{b}W^- \rightarrow b\ell^+\nu\bar{b}\ell^-\bar{\nu}$ are presented. The search is performed in a data sample corresponding to a total integrated luminosity of 4.7 fb^{-1} of pp collisions at a centre-of-mass energy of 7 TeV, collected by the CMS experiment at the LHC. The observed number of events agrees with the standard model prediction, and no evidence of $t'\bar{t}'$ production is found. Upper limits on the production cross section as a function of t' mass are presented. A t' with a mass below $552 \text{ GeV}/c^2$ is excluded at the 95% C.L.



The Compact Muon Solenoid Experiment Analysis Note

The content of this note is intended for CMS internal use and distribution only



12 December 2011 (v5, 17 January 2012)

Search for Heavy Top-like Quark Pair Production
in the Dilepton Final State in pp Collisions at
 $\sqrt{s} = 7$ TeV

D. Barge, C. Campagnari, D. Kovalskyi, V. Krutelyov

University of California, Santa Barbara

W. Andrews, G. Cerati, D. Evans, F. Golf, I. Macneill, S. Padhi, Y. Tu, F. Würthwein, A. Yagil, J. Yoo

University of California, San Diego

L. Bauerdick, K. Burkett, I. Fisk, Y. Gao, O. Gutsche, B. Hooberman, S. Jindariani, J. Linacre, S. Tkaczyk

Fermi National Accelerator Laboratory, Batavia, Illinois

Abstract

The results of a search for pair production of a heavy top-like quark, t' , in the decay mode $t'\bar{t}' \rightarrow bW^+\bar{b}W^- \rightarrow b\ell^+\nu\bar{b}\ell^-\bar{\nu}$ are presented. The search is performed in a data sample corresponding to a total integrated luminosity of 4.7 fb^{-1} of pp collisions at a centre-of-mass energy of 7 TeV, collected by the CMS experiment at the LHC. The observed number of events agrees with the standard model prediction, and no evidence of $t'\bar{t}'$ production is found. Upper limits on the production cross section as a function of t' mass are presented. A t' with a mass below $552 \text{ GeV}/c^2$ is excluded at the 95% C.L.

<http://cms.cern.ch/iCMS/analysisadmin/cadi?ancode=EXO-11-050>

► Thanks to the ARC:


Philip Harris (MIT), Ia Iashvili (Buffalo), Isabel Josa Mutuberria (Madrid, CIEMAT),
Mayda Velasco (Northwestern) (Chairperson)

- ▶ 4th generation of fermions is a simple extension to the SM
- ▶ With a 4th generation, indirect bounds on M_H can be relaxed
- ▶ May possess sufficient intrinsic matter-antimatter asymmetry to be relevant to the baryon asymmetry of the Universe

- ▶ Lepton-Photon result (1.14/fb): $m_{t'} > 422 \text{ GeV}$
 - ▶ <http://cdsweb.cern.ch/record/1376672/files/EXO-11-050-pas.pdf>

- ▶ Tevatron limits already exceeded:
 - ▶ CDF: $m_{t'} > 358 \text{ GeV}$ (<http://arxiv.org/abs/1107.3875>)
 - ▶ D0: $m_{t'} > 285 \text{ GeV}$ (<http://arxiv.org/abs/1104.4522>)

- ▶ We search in the dilepton final state, where
$$t'\bar{t}' \rightarrow W^+bW^-\bar{b} \rightarrow \ell^+\nu b\ell^-\bar{\nu}\bar{b}, \text{ where } \ell = e \text{ or } \ell = \mu.$$



opposite sign dileptons
2 b jets
MET

- ▶ Datasets: DoubleElectron, DoubleMu, MuEG
 - ▶ High p_T dilepton triggers
 - ▶ May10th rereco + Prompt v4 + Aug05th rereco + Prompt v6 + 2011B Data (4.7 fb^{-1})
- ▶ Summer 11 MC

- ▶ Use opposite sign dilepton SUSY analysis lepton selection (SUS-10-007) plus MET cut and b tagging requirement
 - ▶ purpose of this preselection is to reject events other than $t'\bar{t}'$ or $t\bar{t}$, and it is used as a control region for data-MC comparison
- ▶ Define a signal region to separate $t'\bar{t}'$ signal from $t\bar{t}$ background
- ▶ Estimate residual backgrounds using data driven methods where possible
- ▶ Perform the analysis as counting experiment
- ▶ In the absence of an excess, set 95% C.L. upper limits on t' pair production cross sections and lower limit on t' mass



- ▶ Event cleaning: if ≥ 10 tracks; at least 25% purity; at least 1 good DA vertex (not isFake, $\text{ndf} > 4$, $\rho < 2$ cm, $z < 24$ cm)
- ▶ 2 opposite sign isolated leptons with $p_T > 20$ GeV and $|\eta| < 2.5$ (2.4) for e (μ)
- ▶ ≥ 2 pf jets with $p_T > 30$ GeV, $|\eta| < 2.5$
 - ▶ loose pfjet ID (L1FastL2L3 corrected)
 - ▶ $\Delta R > 0.4$ from all leptons passing analysis selection
 - ▶ exactly 2 b tags: TCHM (changed from ≥ 2 in 1.14/fb analysis)
- ▶ MET > 50 GeV (changed from 30 GeV in 1.14/fb analysis)
- ▶ Z veto: $76 < m_{ll} < 106$ GeV veto (for SF leptons)
- ▶ $m_{ll} > 12$ GeV to veto low mass resonances (SF leptons)

Paper and AN




Sample	ee	$\mu\mu$	$e\mu$	all
$t'\bar{t}', M_{t'} = 350 \text{ GeV}/c^2$	22.7 ± 2.0	27.2 ± 2.1	56.1 ± 3.1	106.0 ± 4.2
$t'\bar{t}', M_{t'} = 400 \text{ GeV}/c^2$	10.0 ± 0.9	13.0 ± 1.0	27.7 ± 1.4	50.7 ± 1.9
$t'\bar{t}', M_{t'} = 450 \text{ GeV}/c^2$	5.3 ± 0.4	6.0 ± 0.5	14.3 ± 0.7	25.7 ± 1.0
$t'\bar{t}', M_{t'} = 500 \text{ GeV}/c^2$	2.8 ± 0.2	3.1 ± 0.2	6.3 ± 0.3	12.2 ± 0.5
$t'\bar{t}', M_{t'} = 550 \text{ GeV}/c^2$	1.7 ± 0.1	1.7 ± 0.1	3.4 ± 0.2	6.8 ± 0.3
$t'\bar{t}', M_{t'} = 600 \text{ GeV}/c^2$	0.8 ± 0.1	1.0 ± 0.1	2.1 ± 0.1	3.9 ± 0.2
$t\bar{t} \rightarrow \ell^+ \ell^-$	494.2 ± 11.2	622.3 ± 12.1	1490.7 ± 19.1	2607.2 ± 25.3
$t\bar{t} \rightarrow \text{fake}$	7.3 ± 1.4	0.5 ± 0.3	10.7 ± 1.6	18.4 ± 2.1
$W + \text{jets}$	< 1.8	< 1.8	< 1.8	< 1.8
$\text{DY} \rightarrow \ell^+ \ell^-$	2.7 ± 1.4	1.5 ± 0.9	0.5 ± 0.5	4.8 ± 1.7
Di-boson	0.5 ± 0.1	1.0 ± 0.1	1.8 ± 0.2	3.3 ± 0.3
Single top	14.7 ± 0.9	18.3 ± 1.0	44.1 ± 1.6	77.1 ± 2.1
Total Background	519.4 ± 11.4	643.6 ± 12.2	1547.8 ± 19.3	2710.9 ± 25.5
Data	510	615	1487	2612

Uncertainties are statistical only


- ▶ MC events are weighted to match trigger efficiency, b tagging efficiency, and number of vertices distribution in data
- ▶ 97% of MC background yield is from $t\bar{t}$
- ▶ Data-MC agreement is reasonable
- ▶ Plots on next slide



- ▶ Vertex reweighting has been applied to MC
- ▶ Selected plots: #jets, #vertices, leading lepton and jet p_T , MET
- ▶ More plots in backup
- ▶ Data and MC agreement is reasonable



- ▶ We can discriminate t' from t based on mass
- ▶ For leptonic decay, invariant mass of the lepton and b is constrained as $M_{lb}^2 < M_{t/t'}^2 - M_W^2$



- ▶ We have 2 leptonic decays, so don't know which lepton goes with which b
 - ▶ 4 possible combinations
- ▶ To remove top background, require all 4 $M_{lb} > 170 \text{ GeV}$
 - ▶ equivalently, $M_{lb}^{\min} > 170 \text{ GeV}$
- ▶ Reduces $t\bar{t}$ by 10^4 , and $t't'\bar{} by only \sim 50\%$ (see plot)
- ▶ Very small number of $t\bar{t}$ events still appear in signal region
 - ▶ mostly when at least one of the 2 b s is not from top decay
 - ▶ mistag(s) or extra real b (s)
 - ▶ also some badly mis-measured b jets
- ▶ Signal region definition is changed since 1.14/fb analysis (only 2 of the 4 M_{lb} had to be $> 170 \text{ GeV}$).
 - ▶ requires new data-driven background estimation methods

- ▶ Divide the background events into 4 categories:
 - ▶ Category I: events with mistagged b(s) and real leptons
 - ▶ Category II: events with fake lepton(s) and real bs
 - ▶ Category III: events with 2 real bs and 2 real leptons
 - ▶ Category IV: events with mistagged b(s) and fake lepton(s)
- ▶ Data-driven predictions for Categories I and II using b mistag rate and lepton fake rate (also covers Category IV)
- ▶ No data driven method for Category III: use MC

- ▶ Mistag background: 1 or both the b jets are mistags
- ▶ Estimate from data using method analagous to lepton fake rate, but instead using mistag rate m_i for each jet
 - ▶ Single mistag background contribution, $N_{1\text{-mistag}}$, from events in 1 b-tag bin weighted by $m_i/(1-m_i)$
 - ▶ Double mistag background contribution, $N_{2\text{-mistags}}$, from events in 0 b-tag bin weighted by $m_i/(1-m_i) \times m_j/(1-m_j)$
 - ▶ Final prediction from $N_{1\text{-mistag}} - N_{2\text{-mistags}}$, because $N_{2\text{-mistags}}$ is already counted twice in $N_{1\text{-mistag}}$
- ▶ We see under-prediction of up to 50% in MC closure test
 - ▶ assign 100% systematic uncertainty (but no scaling to avoid over-prediction)
- ▶ Prediction in data: $0.74 \pm 0.27 \pm 0.74$

- ▶ This method is unchanged from the 1.14/fb analysis
- ▶ We estimate the contribution from fake leptons using the data-driven tight-to-loose method described in CMS AN-2010/257
 - ▶ measure tight-to-loose fake rates as a function of lepton p_T and η
 - ▶ estimate number of fakes in data based on number of fakeable objects (FOs)
 - ▶ Expect fake background dominated by $t\bar{t}$ bar decaying to lepton+jets
- ▶ We find zero FOs, thus our fake estimate is 0 events
- ▶ Take upper limit from weight of one FO, giving estimate of $0_{-0}^{+0.4}$
- ▶ Consistent with MC where we also see 0 events with fake leptons

Signal region MC yields:

AN (and paper for 3 t' samples only)

Sample	ee	$\mu\mu$	$e\mu$	all
$t't', M_{t'} = 350 \text{ GeV}/c^2$	5.10 ± 0.96	7.71 ± 1.10	11.35 ± 1.35	24.16 ± 1.98
$t'\bar{t}', M_{t'} = 400 \text{ GeV}/c^2$	3.28 ± 0.48	5.14 ± 0.60	10.49 ± 0.87	18.91 ± 1.16
$t'\bar{t}', M_{t'} = 450 \text{ GeV}/c^2$	2.24 ± 0.29	2.48 ± 0.29	6.17 ± 0.46	10.88 ± 0.62
$t'\bar{t}', M_{t'} = 500 \text{ GeV}/c^2$	1.36 ± 0.16	1.82 ± 0.18	3.15 ± 0.23	6.32 ± 0.33
$t'\bar{t}', M_{t'} = 550 \text{ GeV}/c^2$	0.95 ± 0.11	0.98 ± 0.10	1.92 ± 0.14	3.86 ± 0.21
$t'\bar{t}', M_{t'} = 600 \text{ GeV}/c^2$	0.53 ± 0.06	0.56 ± 0.06	1.25 ± 0.09	2.34 ± 0.12
$t\bar{t} \rightarrow \ell^+\ell^-$	0	0.53 ± 0.34	0	0.53 ± 0.34
$t\bar{t} \rightarrow \text{fake}$	0	0	0	0
$W + \text{jets}$	0	0	0	0
$DY \rightarrow \ell^+\ell^-$	0.66 ± 0.66	0	0	0.66 ± 0.66
Di-boson	0.05 ± 0.04	0.12 ± 0.05	0.12 ± 0.05	0.29 ± 0.08
Single top	0	0.12 ± 0.09	0.36 ± 0.16	0.48 ± 0.18
Total Expected Background	0.71 ± 0.66	0.78 ± 0.36	0.48 ± 0.17	1.97 ± 0.77
Data	0	0	1	1

Very few MC background events pass selection, large stat uncertainties

Category III prediction taken from number of MC events with 2 real leptons and 2 real bs: 0.99 ± 0.69

Summary of background predictions (highlighted predictions are used for total):

Category	Data-driven pred	MC yield
I	$0.74 \pm 0.27 \pm 0.74$	0.98 ± 0.34
II	$0.0^{+0.4}$	0
III	N/A	0.99 ± 0.69
IV	covered by I and II	0
Total prediction	1.73 ± 1.12	

► Uncertainty on Signal Acceptance and Efficiency:

AN

Source	Relative systematic uncertainty
b tagging efficiency	4% (15% for $p_T > 240$ GeV/ c) per b jet
Trigger efficiency	2%
Lepton ID and iso	2% per lepton
Jet and E_T^{miss} energy scale	8%
Total	19%


► Luminosity uncertainty: 4.5%

► Uncertainties on data-driven background estimates dominated by 100% systematic on mistag estimate

- ▶ 1 event observed ($e\mu$ channel)
- ▶ Prediction and observation consistent within the uncertainty
- ▶ No evidence of excess above SM
- ▶ Proceed to set 95% limits on t' pair production cross section

Paper and AN


Sample	Yield
Category I (data-driven)	0.74 ± 0.79
Category II (data-driven)	$0^{+0.4}_{-0.0}$
Category III (simulated)	0.99 ± 0.69
Total prediction	1.73 ± 1.12
Data	1



► Limits evaluated using LandS, taking into account change of signal acceptance with $M_{t'}$

Paper and AN

Paper and AN



Sample	Eff \times Acc \times BR
$t'\bar{t}'$, $M_{t'} = 350$ GeV/c ²	0.16%
$t'\bar{t}'$, $M_{t'} = 400$ GeV/c ²	0.29%
$t'\bar{t}'$, $M_{t'} = 450$ GeV/c ²	0.35%
$t'\bar{t}'$, $M_{t'} = 500$ GeV/c ²	0.41%
$t'\bar{t}'$, $M_{t'} = 550$ GeV/c ²	0.48%
$t'\bar{t}'$, $M_{t'} = 600$ GeV/c ²	0.54%

- Mass limits based on theoretical $t'\bar{t}'$ cross sections (HATHOR)
- **552 GeV observed limit**
- **542 GeV expected limit**

Paper and AN

$M_{t'}$	350 GeV/c ²	400 GeV/c ²	450 GeV/c ²	500 GeV/c ²	550 GeV/c ²	600 GeV/c ²
Theory (pb)	3.200	1.406	0.622	0.330	0.171	0.092
Expected (pb)	0.560	0.309	0.256	0.219	0.187	0.166
Observed (pb)	0.503	0.278	0.230	0.196	0.168	0.149

- ▶ A search for a heavy top-like quark in the dilepton channel is performed with 4.7 fb^{-1} of 2011 data.
- ▶ Data driven estimates are made for mistag and fake backgrounds
- ▶ A MC estimate is used for backgrounds with real bs and leptons
- ▶ We calculate 95% C.L. upper limits on $pp \rightarrow t' t' \bar{t}$ production cross sections as a function of t' mass.
- ▶ Observed lower limit on the t' mass is 552 GeV

▶ Electron selection

- ▶ $p_T > 20 \text{ GeV}; |\eta| < 2.5$
- ▶ VBTF90 (cuts tightened to match Calold+TrklVL HLT requirements)
- ▶ $d_0 \text{ (PV)} < 0.04 \text{ cm}, dz \text{ (PV)} < 1 \text{ cm}$
--calculated w.r.t. 1st good DA PV
- ▶ no muon $\Delta R < 0.1$
- ▶ ≤ 1 miss hits, $|dist| < 0.02 \text{ cm}$ and < 0.02 , CMS AN-2009-159
- ▶ Veto electrons with a supercluster in the transition region ($1.44 < |\eta| < 1.56$)
- ▶ $iso/p_T < 0.15$ (EB pedestal subtraction 1 GeV, no fastjet correction)
- ▶ $ecaliso/p_T < 0.2$

▶ Muon selection

- ▶ $p_T > 20 \text{ GeV}; |\eta| < 2.4$
- ▶ global and tracker muon
- ▶ $\chi^2/ndf < 10$
- ▶ $nValidHits > 10$ -- to be updated to frac of validHits
- ▶ valid StandAloneHits > 0
- ▶ $d_0 \text{ (PV)} < 0.02 \text{ cm}, dz \text{ (PV)} < 1 \text{ cm}$
--calculated w.r.t. 1st good DA PV
- ▶ $(p_T)/p_T < 0.1$
- ▶ $iso/p_T < 0.15$ (no fastjet correction)

- Double Electron
 - HLT_Ele17_CaloIdL_CaloIsoVL_Ele8_CaloIdL_CaloIsoVL
 - HLT_Ele17_CaloIdT_TrkIdVL_CaloIsoVL_TrkIsoVL_Ele8_CaloIdT_TrkIdVL_CaloIsoVL_TrkIsoVL
 - HLT_Ele17_CaloIdT_CaloIsoVL_TrkIdVL_TrkIsoVL_Ele8_CaloIdT_CaloIsoVL_TrkIdVL_TrkIsoVL
- Double Muon
 - HLT_DoubleMu7
 - HLT_Mu13_Mu7
 - HLT_Mu13_Mu8
 - HLT_Mu17_Mu8
- Electron Muon
 - HLT_Mu17_Ele8_CaloIdL
 - HLT_Mu8_Ele17_CaloIdL
 - HLT_Mu17_Ele8_CaloIdT_CaloIsoVL
 - HLT_Mu8_Ele17_CaloIdT_CaloIsoVL

For the high p_T dilepton triggers, the efficiencies listed in Table 1, Table 2, Table 3 and Table 4 are applied to ee , $\mu\mu$ and $e\mu$ Monte Carlo Events. Details of the measurement of the trigger efficiencies are described in [12].

Table 1: The efficiency of the leading leg requirement for the double electron trigger, averaged over the full 2011 data.

Measurement	$0.0 \leq \eta < 1.5$	$1.5 \leq \eta < 2.5$
$20 \leq p_T \leq 30$	0.9849 ± 0.0003	0.9774 ± 0.0007
$p_T > 30$	0.9928 ± 0.0001	0.9938 ± 0.0001

Table 2: The efficiency of the trailing leg requirement for the double electron trigger, averaged over the full 2011 data.

Measurement	$0.0 \leq \eta < 1.5$	$1.5 \leq \eta < 2.5$
$20 \leq p_T \leq 30$	0.9923 ± 0.0002	0.9953 ± 0.0003
$p_T > 30$	0.9948 ± 0.0001	0.9956 ± 0.0001

AN

Table 3: The efficiency of the leading leg requirement for the double muon trigger, averaged over the full 2011 data.

Measurement	$0.0 \leq \eta < 0.8$	$0.8 \leq \eta < 1.2$	$1.2 \leq \eta < 2.1$	$2.1 \leq \eta < 2.4$
$20 \leq p_T \leq 30$	0.9648 ± 0.0007	0.9516 ± 0.0013	0.9480 ± 0.0009	0.8757 ± 0.0026
$p_T > 30$	0.9666 ± 0.0003	0.9521 ± 0.0005	0.9485 ± 0.0004	0.8772 ± 0.0012

Table 4: The efficiency of the trailing leg requirement for the double muon trigger, averaged over the full 2011 data.





Measurement	$0.0 \leq \eta < 0.8$	$0.8 \leq \eta < 1.2$	$1.2 \leq \eta < 2.1$	$2.1 \leq \eta < 2.4$
$20 \leq p_T \leq 30$	0.9655 ± 0.0007	0.9535 ± 0.0013	0.9558 ± 0.0009	0.9031 ± 0.0023
$p_T > 30$	0.9670 ± 0.0003	0.9537 ± 0.0005	0.9530 ± 0.0004	0.8992 ± 0.0011

- TTJets_TuneZ2_7TeV-madgraph-tauola_Summer11-PU_S4_START42_V11-v1 , 165.8 pb
- TTJets_TuneZ2_7TeV-madgraph-tauola_Fall111-PU_S6_START42_V14B-v2 , 165.8 pb
- T_TuneZ2_tW-channel_7TeV-madgraph_Summer11-PU_S4_START42_V11-v1 , 7.87 pb
- T_TuneZ2_t-channel_7TeV-madgraph_Summer11-PU_S4_START42_V11-v1 , 41.92 pb
- T_TuneZ2_s-channel_7TeV-madgraph_Summer11-PU_S4_START42_V11-v1 , 3.19 pb
- Tbar_TuneZ2_tW-channel_7TeV-madgraph_Summer11-PU_S4_START42_V11-v1 , 7.87 pb
- Tbar_TuneZ2_t-channel_7TeV-madgraph_Summer11-PU_S4_START42_V11-v1 , 22.65 pb
- Tbar_TuneZ2_s-channel_7TeV-madgraph_Summer11-PU_S4_START42_V11-v1 , 1.44 pb
- WJetsToLNu_TuneZ2_7TeV-madgraph-tauola_Summer11-PU_S4_START42_V11-v1 , 31314 pb
- DYJetsToLL_TuneD6T_M-50_7TeV-madgraph-tauola_Summer11-PU_S4_START42_V11-v1 , 3048 pb
- DYToEE_M-20_CT10_TuneZ2_7TeV-powheg-pythia_Summer11-PU_S4_START42_V11-v1 , 1666 pb
- DYToMuMu_M-20_CT10_TuneZ2_7TeV-powheg-pythia_Summer11-PU_S4_START42_V11-v1 , 1666 pb
- DYToTauTau_M-20_CT10_TuneZ2_7TeV-powheg-pythia-tauola_Summer11-PU_S4_START42_V11-v1 , 1666 pb
- DYToEE_M-10To20_TuneZ2_7TeV-pythia6_Summer11-PU_S4_START42_V11-v1 , 3319.61 pb
- DYToMuMu_M-10To20_TuneZ2_7TeV-pythia6_Summer11-PU_S4_START42_V11-v1 , 3319.61 pb
- DYToTauTau_M-10To20_CT10_TuneZ2_7TeV-powheg-pythia-tauola_Summer11-PU_S4_START42_V11-v2 , 3319.61 pb
- WWJetsTo2L2Nu_TuneZ2_7TeV-madgraph-tauola_ummer11-PU_S4_START42_V11-v1, 4.783 pb
- WZJetsTo2L2Q_TuneZ2_7TeV-madgraph-tauola_Summer11-PU_S4_START42_V11-v1, 1.786 pb
- WZJetsTo3LNu_TuneZ2_7TeV-madgraph-tauola_Summer11-PU_S4_START42_V11-v1, 0.856 pb
- ZZJetsTo2L2Nu_TuneZ2_7TeV-madgraph-tauola_Summer11-PU_S4_START42_V11-v1, 0.30 pb
- ZZJetsTo2L2Q_TuneZ2_7TeV-madgraph-tauola_Summer11-PU_S4_START42_V11-v1, 1.0 pb
- ZZJetsTo4L_TuneZ2_7TeV-madgraph-tauola/_ummer11-PU_S4_START42_V11-v1, 0.076 pb
- TprimeTprimeToBWBWinc_M-350_7TeV-madgraph_Summer11-PU_S4_START42_V11-v2, 3.200 pb
- TprimeTprimeToBWBWinc_M-400_7TeV-madgraph_Summer11-PU_S4_START42_V11-v2, 1.406 pb
- TprimeTprimeToBWBWinc_M-450_7TeV-madgraph_Summer11-PU_S4_START42_V11-v2, 0.622 pb
- TprimeTprimeToBWBWinc_M-500_7TeV-madgraph_Summer11-PU_S4_START42_V11-v2, 0.330 pb
- TprimeTprimeToBWBWinc_M-550_7TeV-madgraph_Summer11-PU_S4_START42_V11-v2, 0.171 pb
- TprimeTprimeToBWBWinc_M-600_7TeV-madgraph_Summer11-PU_S4_START42_V11-v2, 0.0923 pb

Mistag method: mistag rate

► Mistag rate is taken from CMS PAS BTV-11-001

<http://cms-physics.web.cern.ch/cms-physics/public/BTV-11-001-pas.pdf>



- ▶ Prediction of the method compared to true count of events with mistags in MC
- ▶ Total mistags predicted: 1.30 ± 0.24 (uncertainties stat only)
- ▶ Total mistags actual: 0.98 ± 0.34

- ▶ Repeat using Fall I I ttbar sample (20 times larger)
 - ▶ predicted (ttbar only): 0.27 ± 0.02
 - ▶ observed (ttbar only): 0.45 ± 0.09

- ▶ Repeat for preselection region
 - ▶ predicted: 155.2 ± 1.6
 - ▶ observed: 288.3 ± 9.2